

Split Prep Instructions for CLENPIQ®

PREPARATION FOR COLONOSCOPY

(Follow ONLY these instructions given by the GI Department)



READ ALL PREP INSTRUCTIONS.

- WEIGHT LOSS OR DIABETES MEDICATIONS THAT SLOW DIGESTION MAY NEED TO BE STOPPED. Please speak with our department about instructions to hold if you are taking these medications.
- ASK YOUR PRESCRIBING DOCTOR ABOUT WHEN TO STOP TAKING BLOOD THINNERS such as Plavix, Xarelto, Eliquis, Coumadin, or Brulinta
- IF YOU TAKE INSULIN, ask your prescribing doctor about dosage for your preparation day. Do not take any insulin the day of the procedure.
- STOP TAKING all herbal products, oil-based vitamins, and iron supplements.
- STOP TAKING fiber supplements such as Metamucil, Citrucel, FiberCon, BeneFiber, or Konsyl.
- STOP TAKING medicine that stops diarrhea such as Imodium, Kaopectate, or Pepto Bismol.
- STOP EATING salads, nuts, seeds, popcorn, hard fruits, and hard vegetables.

MAKE SURE SOMEONE IS AVAILABLE TO DRIVE YOU HOME AFTER YOUR PROCEDURE.



GET THESE ITEMS FROM THE STORE AND PHARMACY:

- CLENPIQ BOWEL PREP from pharmacy.
- CLEAR LIQUIDS TO DRINK FOR THE DAY BEFORE YOUR COLONOSCOPY (no red or purple). You can find a list of approved clear liquids on your patient checklist.

FOR PATIENTS WITH DIABETES OR KIDNEY DISEASE:

USE WATER, GATORADE ZERO, OR CRYSTAL LIGHT INSTEAD OF REGULAR GATORADE.

Examples of clear liquids: (NO RED OR PURPLE COLOR)

- Broth (clear only)
- Clear hard candy
- Coffee (no milk)
- Jell-0
- Juice (clear, no pulp)
- Kool-Aid
- Popsicle
- Soda
- Tea

THE DAY BEFORE YOUR COLONOSCOPY

- DRINK ONLY CLEAR LIQUIDS ALL DAY
- NO SOLID FOOD
- NO ALCOHOL

- DRINK AT LEAST EIGHT (8) OUNCES OF CLEAR LIQUIDS EVERY HOUR while awake.
- WAIT EIGHT (8) HOURS BETWEEN YOUR FIRST AND SECOND DOSE OF THIS PREP.
- O AT 4 P.M., DRINK ONE (1) ENTIRE BOTTLE OF CLENPIQ.
- AFTER DRINKING THE CLENPIQ, YOU MUST DRINK FIVE (5) 8-OZ GLASSES OF CLEAR LIQUIDS over the next five (5) hours.

YOU WILL GET A CALL ON THIS DAY ABOUT WHEN TO COME IN FOR YOUR PROCEDURE.



- NO SOLID FOOD
- NO ALCOHOL
- NO GUM OR MINTS
- FIVE (5) HOURS BEFORE
- YOUR PROCEDURE ARRIVAL TIME, DRINK ANOTHER ONE (1) BOTTLE OF CLENPIQ. You must finish the entire bottle.
- O AFTER DRINKING THE CLENPIQ, YOU MUST DRINK THREE (3) 8-0Z GLASSES OF CLEAR LIQUIDS over the next 1-2 hours.
- STOP DRINKING ALL LIQUIDS THREE (3) HOURS BEFORE YOUR ARRIVAL TIME. If you don't, your procedure may be delayed or canceled.
- Take your morning asthma and seizure medications with a small sip of water.
- If you take medicine for your heart or blood pressure, please talk to our nurse before your procedure.

AFTER YOUR PROCEDURE, YOU MAY EAT YOUR USUAL DIET UNLESS YOUR DOCTOR TELLS YOU DIFFERENTLY.

THIS HELPS CLEAN YOUR BOWEL WELL, EVEN

IF YOU'RE ALREADY PASSING CLEAR LIQUID.

> YOU NEED TO DRINK ALL OF THE PREPARATION SOLUTION AS INSTRUCTED!

If you think the prep is not working by 10 P.M., call 856.342.2000 and ask for the Gastroenterology Fellow on call.

IF YOU HAVE ANY QUESTIONS OR CONCERNS, DON'T HESITATE TO CALL.

(MONDAY-FRIDAY, 8 A.M.-4:30 P.M.) Please call the location you are scheduled to have your procedure.

• CAMDEN: 856.342.2047

- MOORESTOWN: 856.533.1961
- MOUNT LAUREL: 856.380.7750
- SALEM: 856.339.6040
- WILLINGBORO: 609.589.3651



2840/6-24/HM